

16. Ochert, A. S., Boulter, A. W., Birnbaum, W., Johnson, N. W., and Teo, C. G. (1994) Inhibitory effect of salivary fluids on PCR: potency and removal. *PCR Methods Applic.* **3**, 365–368.
17. Felipe, M. S. S., Rogelin, R., Azevedo, M. O., and Astolfifilho, S. (1993) Extended application of an efficient method for RNA isolation from different organisms. *Biotechniques* **7**, 639–644.
18. Wan, C. Y. and Wilkins, T. A. (1994) A modified hot borate method significantly enhances the yield of high-quality RNA from cotton (*Gossypium hirsutum* L). *Anal. Biochem.* **223**, 7–12.
19. Manning, K. (1991) Isolation of nucleic acids from plants by differential solvent precipitation. *Anal. Biochem.* **195**, 45–50.
20. Lodhi, M. A., Ye, G.-N., Weeden, N. F., and Reisch, B. I. (1994) A simple and efficient method for DNA extraction from grapevine cultivars and *Vitis* species. *Plant Mol. Biol. Reporter* **12**, 6–13.
21. Poulson R. (1973) Isolation, purification and fractionation of RNA, in *The Ribonucleic Acids* (Stewart, P. R. and Letham, D. S.), Springer-Verlag, Berlin, pp. 243–261.
22. Kwok, S. and Higuchi, R. (1989). Avoiding false positives with PCR. *Nature* **339**, 237,238.
23. Winn-Deen, E. S., Batt, C. A., and Wiedmann, M. (1993) Non-radioactive detection of *Mycobacterium tuberculosis* LCR products in a microtitre plate format. *Mol. Cell. Probes* **7**, 179–186.
24. Ferre, F. (1992) Quantitative or semi-quantative PCR: reality versus myth. *PCR Methods Applic.* **2**, 1–9.